

Decomposition theorem for invertible substitutions on three-letter alphabet *

Bo TAN[†] Zhi-Xiong WEN[†] Yiping ZHANG[†]

Abstract

We study the structure of invertible substitutions on three-letter alphabet. We show that there exists a finite set $\mathbb{S}$ of invertible substitutions such that any invertible substitution can be written as $I_w \circ \sigma_1 \circ \sigma_2 \circ \cdots \circ \sigma_k$, where I_w is the inner automorphism associated with w , and $\sigma_j \in \mathbb{S}$ for $1 \leq j \leq k$. As a consequence, M is the matrix of an invertible substitution if and only if it is a finite product of non-negative elementary matrices.

Keywords: Invertible substitution, Indecomposable substitution, Inner automorphism

2000 Mathematics Subject Classification: Primary 20M05; Secondary 68R15.

1 Introduction

The study of substitutions (endomorphisms of the free monoid of finite type) plays an important role in finite automata, symbolic dynamics, and fractal geometry ([1, 3, 7, 18, 21]). It has various applications in quasicrystals, computational complexity, information theory (see for instance [2, 4, 6, 10]). In addition, substitution is also a fundamental object studied in combinatorial group theory [9, 11, 12].

For substitutions over two-letter alphabet plenty of results have been obtained [5, 13, 20, 24]. The notion of invertible substitution appears in [16]: these are the substitutions which extend as an automorphism of the corresponding free group. Since then, they have been studied by many authors (see for example [5, 8, 23]).

The invertible substitutions over a two-letter alphabet form a monoid whose structure is known [22]: this monoid can be generated by a permutation and two so-called Fibonacci substitutions (see Theorem 2.1 below).

*Research supported by NSFC and by the Special Funds for Major State Basic Research Projects of China

[†]Department of Mathematics, Wuhan University, 430072 Hubei, Wuhan, P. R. China, e-mail: tanbo@colmath.whu.edu.cn, zhixwen@whu.edu.cn, ypzhang@whu.edu.cn

This result on the structure of invertible substitutions had many applications, namely to the study of local isomorphisms of fixed points of substitutions [22, 25] and to the study of trace maps [16].

When the alphabet has more than two letters, the situation is much more complicated. In [26], it was shown, by enumerating infinitely many so-called indecomposable substitutions, that the monoid of invertible substitutions over three letters (which will be denoted by $\mathcal{IS}(A^*)$) is not finitely generated. So, up to now the structure of $\mathcal{IS}(A^*)$ remained unknown.

In this paper, we elucidate the structure of $\mathcal{IS}(A^*)$. We show that if σ is an invertible substitution over a three-letter alphabet, there exists a word w such that $I_w \circ \sigma$ or $I_{w^{-1}} \circ \sigma$ is the composition of finitely many Fibonacci substitutions and permutations (Theorem 3.1). As a consequence, the substitution matrix of an invertible substitution is positively decomposable (Theorem 3.2).

2 Preliminaries and notations

Let us first recall some basic definitions and notations in the theory of substitutions (see [1, 18, 12] for a general theory).

Let $A = \{a, b, c\}$ (resp. $\bar{A} = \{a, b, c, a^{-1}, b^{-1}, c^{-1}\}$), A^* (resp. Γ_A) be the free monoid (resp. the free group) generated by A (the unit element is the empty word ε). The elements in A^* will be called “positive words” or simply “words” and those of Γ_A “signed” or “mixed” words. The inverse of a positive word will be said to be “negative”.

Let $w \in \Gamma_A$. If $w = x_1 \cdots x_k$ with $x_i \in \bar{A}$ ($i = 1, 2, \dots, k$) and $x_i x_{i+1} \neq \varepsilon$ ($i = 1, \dots, k-1$), we say that $x_1 \cdots x_k$ is in the reduced form and that the length of w is k . This length will be denoted by $|w|$.

Let w, w_j ($j = 1, \dots, k$) $\in \Gamma_A$. If $w = w_1 w_2 \cdots w_k$ satisfies $|w| = |w_1| + \cdots + |w_k|$, we say that $w_1 w_2 \cdots w_k$ is a reduced expression of w . We then say that w_1 is a prefix of w and that w_k is a suffix of w , and we then write $w_1 \triangleleft w$ and $w_k \triangleright w$ respectively.

As already defined, a substitution over A is a morphism σ of A^* . Such a morphism extends in a natural way to an endomorphism of Γ_A ; If this extension is an automorphism of Γ_A , the substitution σ is said to be invertible. The set of substitutions (resp. invertible substitutions) is denoted by $\mathcal{S}(A^*)$ (resp. by $\mathcal{IS}(A^*)$).

We often identify an endomorphism σ of Γ_A with the triple $(\sigma(a), \sigma(b), \sigma(c))$ of (maybe mixed) words. We define the length of σ to be $|\sigma| = |\sigma(a)| + |\sigma(b)| + |\sigma(c)|$.

If U is a subset of a monoid, $\langle U \rangle$ stands for the sub-monoid (not the sub-group, even when dealing inside a group) generated by U . The use of the same notation for different groups and monoids will not generate any confusion.

We shall use the following basic invertible substitutions and automorphisms.

• Permutations

$\mathcal{P}$ will denote the symmetric group on A . Notice that $\mathcal{P} = \langle \pi_1, \pi_2 \rangle$, where $\pi_1 = (b, a, c)$, $\pi_2 = (c, b, a)$. Note that the identity $I = (a, b, c) \in \mathcal{P}$.

- **Fibonacci type:**

Set $\phi_l = (ba, b, c)$, $\phi_r = (ab, b, c)$.

Set $\mathbf{L} = \{\pi \circ \phi_l \circ \pi'; \pi, \pi' \in \mathcal{P}\}$, $\mathcal{R} = \{\pi \circ \phi_r \circ \pi'; \pi, \pi' \in \mathcal{P}\}$, and $\mathcal{F} = \mathcal{R} \cup \mathbf{L}$.

The elements in $\mathcal{F}$ are called substitutions of Fibonacci type or simply Fibonacci substitutions.

- **Simple substitutions**

It is easy to check the following equalities.

$$S := \langle \pi_1, \pi_2, \phi_l, \phi_r \rangle = \langle \mathcal{P}, \mathbf{L}, \mathcal{R} \rangle \subset \mathcal{IS}(A^*)$$

Definition 2.1. *The elements in S will be called simple substitutions.*

- **Involutions**

$$\iota_1 := (a^{-1}, b, c), \iota_2 := (a, b^{-1}, c), \iota_3 := (a, b, c^{-1}), \mathcal{I} := \{\iota_i ; 1 \leq i \leq 3\}.$$

It is well known (the cancellation theory of Nielsen, see for example [12]) that

$$\text{Aut}(\Gamma_A) = \langle \mathcal{P}, \mathcal{F}, \mathcal{I} \rangle = \langle \pi_1, \pi_2, \phi_l, \phi_r, \iota_1 \rangle.$$

For the case $A = \{a, b\}$, the following decomposition is known.

Theorem 2.1. [22] *Let $A_2 = \{a, b\}$, $\alpha = (b, a)$, $\beta = (ab, b)$, $\gamma = (ba, b)$. Then we have*

$$\mathcal{IS}(A_2^*) = \langle \alpha, \beta, \gamma \rangle. \quad (2.1)$$

Remark 2.1. *With our terminology, the above theorem says that, for the case of two-letter alphabet, every invertible substitution is simple. In the case $A = \{a, b, c\}$ it is shown in [26] that there exist non-simple substitutions and that there exists no finite set of substitutions which generates $\mathcal{IS}(A^*)$. Consequently the above theorem cannot be extended to the case of three letters. Our Theorem 3.1 below will give a characterization of the structure of $\mathcal{IS}(A^*)$.*

- **Indecomposable substitutions**

Definition 2.2. *Let σ be an invertible substitution.*

σ is called trivial (resp. non-trivial) if $\sigma \in \mathcal{P}$ (resp. $\sigma \notin \mathcal{P}$).

If there exist non-trivial invertible substitutions σ_1 and σ_2 such that $\sigma = \sigma_1 \circ \sigma_2$, we say that σ is decomposable. Otherwise, we say that σ is indecomposable.

The elements in $\mathcal{P} \cup \mathcal{F}$ are indecomposable in the sense of this definition. They are “simple indecomposable substitutions” according to Definition 2.1. The others will be called non-simple indecomposable substitutions. As a matter of fact, there exist infinitely many indecomposable substitutions, hence $\mathcal{IS}(A^*)$ is not finitely generated [26].

Example 2.1. *Here are some examples of non-simple indecomposable substitutions.*

$$\begin{aligned} &(ab, acb, ac^n) \ (n \geq 2) \\ &(ab^{n+1}c, ab^nc, acc) \ (n \geq 1) \\ &(acbc^n, ac^{n+1}, ab^kc^n) \ (n, k \geq 1) \end{aligned}$$

Notice that any invertible substitution can be written as the product of finitely many indecomposable substitutions.

• Inner automorphisms

Let $z \in \Gamma_A$. $I_z \in \text{Aut}(\Gamma_A)$ is defined as follows: $I_z(w) = zwz^{-1}$ ($w \in \Gamma_A$). That is $I_z = (zaz^{-1}, zbz^{-1}, zcz^{-1})$. We have $I_z \circ (w_1, w_2, w_3) = (zw_1z^{-1}, zw_2z^{-1}, zw_3z^{-1})$. Notice also that $I_\varepsilon = I$.

We shall also use the following operation.

• Cyclic operators:

Let $w = x_1x_2 \cdots x_k \in A^*$. Then for $1 \leq i \leq k$,

$$\begin{aligned} I_{(x_1 \cdots x_i)^{-1}}(w) &: = x_{i+1} \cdots x_k x_1 \cdots x_i \\ I_{x_{i+1} \cdots x_n}(w) &: = x_{i+1} \cdots x_k x_1 \cdots x_i \end{aligned}$$

We observe that the above operator is in fact a k -cyclic operator.

The following operations on substitutions will be frequently used in this paper:

$$\begin{aligned} I_{z^{-1}} \circ (zw_1, zw_2, zw_3) &= (w_1z, w_2z, w_3z), \\ I_z \circ (w_1z, w_2z, w_3z) &= (zw_1, zw_2, zw_3). \end{aligned}$$

More generally, let $\sigma \in \mathcal{S}(A^*)$, $z \in A^*$ (or $z^{-1} \in A^*$) be such that $\sigma' = I_z \circ \sigma$ is still a substitution, we shall say that I_z is a cyclic operator for σ .

• Substitution matrix:

Let $w \in \Gamma_A$. $|w|_a$ (resp. $|w|_b$, $|w|_c$) will denote the number (algebraic sum) of appearances of a (resp. b , c) in w (e.g. $|a^{-1}ba|_a = 0$).

Let $\sigma \in \mathcal{S}(A^*)$. The substitution matrix of σ is defined by:

$$M_\sigma = \begin{pmatrix} |\sigma(a)|_a & |\sigma(b)|_a & |\sigma(c)|_a \\ |\sigma(a)|_b & |\sigma(b)|_b & |\sigma(c)|_b \\ |\sigma(a)|_c & |\sigma(b)|_c & |\sigma(c)|_c \end{pmatrix}$$

Let $\sigma, \tau \in \mathcal{S}(A^*)$. We have $M_{\sigma \circ \tau} = M_\sigma M_\tau$. If $\sigma \in \mathcal{IS}(A^*)$, then $\det(M_\sigma) = \pm 1$.

The above definitions and equalities can be extended to the case where σ and τ are endomorphisms on Γ_A .

Substitution matrix is a basic tool to study substitutions [18], but it cannot characterize the non-commutative combinatorial properties of A^* and $\mathcal{S}(A^*)$.

Notice that the usual non-negative elementary matrices are exactly the substitution matrices of permutations or Fibonacci substitutions.

As have been pointed out in [26], comparing the case of two letters with that of three letters, we have

Remark 2.2. 1) *If $M = (m_{ij})$, $(m_{ij} \in \{0, 1, 2, \dots\})$ is a 2×2 -matrix with determinant $\det(M) = \pm 1$, then M is a finite product of non-negative elementary matrices. In particular, any such matrix is the substitution matrix of some invertible substitution.*

2) *The above statement is no longer true for 3×3 -matrices.*

A simple counter example is [26]

$$M = \begin{pmatrix} 3 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

We can verify that $\det(M) = 1$, that M can not be decomposed as a product of non-negative elementary matrices, and that it is not a substitution matrix of any invertible substitution.

We mention that there are infinitely many such matrices.

It was an open problem whether the substitution matrix of an invertible substitution is a finite product of non-negative elementary matrices [1]. Our Theorem 3.2 will give an affirmative answer.

3 Main theorems

The following decomposition theorem characterizes the structure of $\mathcal{IS}(A^*)$: any invertible substitution is a simple one up to a cyclic operator.

Theorem 3.1. *Let $\sigma \in \mathcal{IS}(A^*)$. There exists $w \in A^*$ or $w^{-1} \in A^*$ such that*

- (1) *$I_w \circ \sigma$ is a simple substitution. In other words, after a cyclic operation $(I_w \circ \sigma)$, any invertible substitution becomes simple.*
- (2) *Furthermore, we can take w (or w^{-1}) to be a common suffix (or prefix) of $\sigma(a)$, $\sigma(b)$ and $\sigma(c)$.*

The following examples illustrate the decomposition after a cyclic operator:

Example 3.1.

1. $\sigma = (ab, acb, ac^2)$. It can be checked that σ is an indecomposable substitution. However we have, by a cyclic operation, $I_{a^{-1}} \circ \sigma = (ba, cba, c^2a) = (b, c, a) \circ (ac, b, c) \circ (a, b, bc)^2 \circ (a, ba, c)$, which is clearly simple.
2. $\sigma = (acbab, acbaccab, accaccab)$ is indecomposable. It is easy to check that $I_b \circ \sigma$ is simple. Notice that in this example, $I_{a^{-1}} \circ \sigma$ is not simple, though it is a substitution.

Remark 3.1. We point out that Theorem 2.1 is an easy consequence of Theorem 3.1.

In fact, let (w_1, w_2) be an invertible substitution, $w_1, w_2 \in \{a, b\}^*$. Then it is obvious that (w_1, w_2, c) is an invertible substitution over $\{a, b, c\}$. Applying Theorem 3.1, and noticing that w in the theorem must be empty, Theorem 2.1 follows.

On the other hand, since $I_{b^{-1}} \circ (ba, b) = (ab, b)$, Theorem 2.1 can be restated as the same as Theorem 3.1:

Any invertible substitution over $\{a, b\}$, having been transformed by some cyclic operator, can be expressed as a finite composition of the Fibonacci (ab, b) and the permutation (b, a) .

Remark 3.2. It should be noticed that Theorem 3.1 can not be extended directly to the case of an alphabet of 4 letters or more.

It suffices to see the following simple example of substitution on $\{a, b, c, d\}$: $\sigma = (ab, acb, acc, d)$. It is indecomposable, but no inner automorphism “ I_z ” can be applied.

Applying the above theorem to the substitution matrices we shall get the following theorem which explains the Remark 2.2:

Theorem 3.2. Let M be a 3×3 -matrix of non-negative integral coefficients. M is the substitution matrix of some invertible substitution if and only if it is a finite product of non-negative elementary matrices.

4 Proofs

The theorems will be proved by several lemmas.

We shall use the following notations. The symbol “+” (resp. “−”) will represent various non-empty positive words (resp. non-empty negative words), which are the elements of $\langle a, b, c \rangle$ (resp. $\langle a^{-1}, b^{-1}, c^{-1} \rangle$). We also use the symbols like “+ − +” to represent various types of mixed words. As an example, $u = + - +$ means that $u = u_1 u_2^{-1} u_3$, where $u_j \in A^*$ and $u_1 u_2^{-1} u_3$ is the reduced expression (i.e. $|u_1 u_2^{-1} u_3| = |u_1| + |u_2| + |u_3|$). The meanings of “ $u = + \dots$ ”, “ $u = \dots -$ ”, “ $u = + \dots -$ ” etc. are now clear.

Recall that by Nielsen's cancellation method, a morphism $\sigma = (w_1, w_2, w_3)$ on Γ_A is invertible if and only if (w_1, w_2, w_3) can be carried by finitely many "elementary Nielsen transformations" into the identity, and each of these elementary Nielsen transformations satisfies some "cancellation" conditions (see [12]). The following lemma is a simple version of Nielsen's cancellation procedure that we use in the proofs. For the details, we refer the reader to [12, 14, 15].

Lemma 4.1. *Let $\sigma = (w_1, w_2, w_3) \in \text{Aut}(\Gamma_A)$. There exist $k \geq 0$ and $\tau_1, \dots, \tau_k \in \{\pi_1, \pi_2, \phi_l, \phi_r, \iota_1\}$ such that, denoting*

$$\sigma_0 = \sigma, \quad \sigma_i = \sigma_{i-1} \circ \tau_i \quad (i = 1, \dots, k), \quad (4.1)$$

$$|\sigma_i| \leq |\sigma_{i-1}| \quad (i = 1, \dots, k) \quad (4.2)$$

and that σ_k is the identity.

Notation: We write the procedure in the above lemma as

$$\sigma \rightarrow \sigma_1 \rightarrow \dots \rightarrow \sigma_k.$$

We say that each arrow represents a "Nielsen's cancellation" and that σ is cancelled to σ_k .

Definition 4.1. *We say that a non-trivial substitution $\sigma = (w_1, w_2, w_3)$ is mixed if it satisfies $w_i w_k^{-1} w_j = + - + \forall i, j, k \in \{1, 2, 3\}, i \neq k, j \neq k$.*

The following lemma is proved in [26].

Lemma 4.2. *Any mixed substitution is non-invertible.*

Corollary 4.1. *Let $\sigma = (w_1, w_2, w_3)$ be an invertible substitution, suppose that for all $i \neq j$, w_i is neither a prefix nor a suffix of w_j . There exist $\pi \in \mathcal{P}$, non-empty words u, v, x, y such that either $\sigma \circ \pi = (ux, uv, yv)$ or $\sigma \circ \pi = (uxv, uv, y)$.*

Proof. By the above lemma, σ is not mixed. Then by the assumption of the lemma we have that, there exist i, j, k ($i \neq j, j \neq k$) such that $w_i^{-1} w_j w_k^{-1} = -$.

Ignoring some permutation, we can only consider the following two possibilities:

1. $w_1^{-1} w_2 w_3^{-1} = -$; 2. $w_1^{-1} w_2 w_1^{-1} = -$.

Case 1: We get $w_2 = uv$ and $w_1 = ux, w_3 = yv$.

Case 2: We must have $w_2 = uv$ and $w_1 = uf = gv$. It's clear that $|u| \neq |g|$, otherwise $u = g$ then $w_1 = w_2$ which contradicts the invertibility.

Subcase 2.1: $|g| > |u|$, then by $w_1 = uf = gv$, we have $g = ux$ and $w_1 = uxv$. Then $\sigma = (uxv, uv, y)$ with $y = w_3$.

Subcase 2.2: $|g| < |u|$, then still by $w_1 = uf = gv$ we have

$u = gx$ and $v = xf$, thus $w_1 = uf$, $w_2 = uxf$, $\sigma = (uf, uxf, y)$. $\square$

Lemma 4.3. *Let $w, v \in \Gamma_A$ and σ be a morphism on Γ_A . We have*

$$I_w \circ I_v = I_{wv}, \quad (I_w)^{-1} = I_{w^{-1}}, \quad \sigma \circ I_w = I_{\sigma(w)} \circ \sigma.$$

Proof. By direct verifications. Let us show the last equality:

$$\text{Let } u \in \Gamma_A. \text{ We have } \sigma \circ I_w(u) = \sigma(wuw^{-1}) = \sigma(w)\sigma(u)\sigma(w)^{-1} = I_{\sigma(w)}(\sigma(u)). \quad \square$$

Let $w = \alpha_1\alpha_2 \cdots \alpha_{|w|} \in \Gamma_A$ ($\alpha_i \in \bar{A}$). We shall use the following notations:

$$h_i(w) = \alpha_i \quad (i = 1, 2, \dots, |w|), \quad h_\infty(w) = h_{|w|}(w). \quad (4.3)$$

Lemma 4.4. *Let $u \in \Gamma_A$ and $x, y \in A^*$ be non-empty. Suppose that*

$$ux = yu \quad (4.4)$$

Then we have either $u \in A^$ or $u^{-1} \in A^*$.*

Proof. First assume that $h_1(u) = +$. We will show that $u \in A^*$.

Otherwise, we may write $u = u_1\beta^{-1}u_2$ (the reduced expression) where $u_1 \in A^*$, $u_1 \neq \epsilon$, $\beta \in A = \{a, b, c\}$ and $u_2 \in \Gamma_A$.

Then by (4.4), we have

$$u_1\beta^{-1}u_2x = yu_1\beta^{-1}u_2 \quad (4.5)$$

Notice that the right hand of the above equation is a reduced expression, and

$$h_{|u_1|+1}(yu_1\beta^{-1}u_2) = +,$$

which, together with (4.5), implies

$$\beta^{-1}u_2x = + \cdots. \quad (4.6)$$

Since $\beta^{-1}u_2$ is a reduced expression and $x \in A^*$, (4.6) implies that $\beta^{-1}u_2x = +$. Returning to (4.5), we have that the left hand of (4.5) is a positive word, which contradicts the right hand.

For the case $h_1(u) = -$, we can show similarly that $u^{-1} \in A^*$. $\square$

Corollary 4.2. *Let $w \in \Gamma_A$, $\sigma = (w_1, w_2, w_3)$ and $\tau = (u_1, u_2, u_3)$ be two substitutions. If $I_w \circ \sigma = \tau$, we have $|w_i| = |u_i|$ ($i = 1, 2, 3$). In particular $|\sigma| = |\tau|$.*

Proof. We have $w^{-1}w_iw = u_i$ ($i = 1, 2, 3$). By the above lemma, we have $w \in A^*$ or $w^{-1} \in A^*$. If $w \in A^*$, we have $w_iw = wu_i$ which implies $|w_i| = |u_i|$; if $w' = w^{-1} \in A^*$, we have $w'w_i = u_iw'$. $\square$

Lemma 4.5. *Let $u, v, x, y \in A^*$. We have*

- (1) *If u is not a suffix of ux , then u is not a suffix of ux^m ($m \geq 1$).*
- (2) *If v is not a prefix of yv , then v is not a prefix of y^mv ($m \geq 1$).*

Proof. We only show that $u \triangleright ux^m$ implies $u \triangleright ux$ by induction on the length of u :

If $|u| \leq |x|$, the conclusion is obvious. Suppose now $|u| > |x|$. Since $u \triangleright ux^m$, we can write $u = u'x$ for some $u' \in A^*$, then $u'x \triangleright u'xx^m$, that is, $u' \triangleright u'x^m$. Since $|u'| < |u|$, by the hypothesis of induction we obtain $u' \triangleright u'x$, which implies $u \triangleright ux$. $\square$

The following lemmas study the “cancellation properties” between special words, where we shall use $h_1(w)$ and $h_\infty(w)$ to indicate the first and the last character of w as defined in (4.3).

Lemma 4.6. *Let u, v, x and y be non-empty words satisfying*

$$h_1(u) \neq h_1(y) \text{ and } h_\infty(x) \neq h_\infty(v); \quad (4.7)$$

$$x \text{ (resp. } v) \text{ is not a prefix of } v \text{ (resp. } x); \quad (4.8)$$

$$y \text{ (resp. } u) \text{ is not a suffix of } u \text{ (resp. } y). \quad (4.9)$$

Let $w_1 = ux, w_2 = uv, w_3 = yv$ and consider the following mixed word:

$$w = w_{i_0}^\epsilon w_{i_1}^{-\epsilon} \cdots w_{i_k}^{(-1)^k \epsilon} \quad (4.10)$$

where $k \geq 0$, $\epsilon \in \{+1, -1\}$, $i_m \in \{1, 2, 3\}$ ($m = 0, 1, \dots, k$) and $i_m \neq i_{m+1}$ ($m = 0, 1, \dots, k-1$). Then we have

$$h_1(w) = h_1(w_{i_0}^\epsilon), \quad h_\infty(w) = h_\infty(w_{i_k}^{(-1)^k \epsilon}) \quad (4.11)$$

and

$$|w| \geq 2. \quad (4.12)$$

(Notice that (4.10) is in general not a reduced expression, this means that there are possible cancellations between words).

Proof. First notice the following obvious facts:

Let $u_1, u_2 \in \Gamma_A^*$ be two (mixed) words in reduced expression. $|u_1 u_2| < |u_1| + |u_2|$ (i.e. some cancellation occurs) if and only if $h_\infty(u_1) = (h_1(u_2))^{-1}$.

If $h_\infty(u_1) \neq (h_1(u_2))^{-1}$, then $v = u_1 u_2$ is a reduced expression, thus $h_1(v) = h_1(u_1)$ and $h_\infty(v) = h_\infty(u_2)$.

We only show $h_1(w) = h_1(w_{i_0}^\epsilon)$. The proof for $h_\infty(w) = h_\infty(w_{i_k}^{(-1)^k \epsilon})$ is similar.

We consider only the case $\epsilon = +1$, the case $\epsilon = -1$ is similar.

When $k \leq 3$, we can check the conclusion directly by enumerating all possible cases. Now we will show it for $k \geq 4$ by induction.

We have to consider carefully several cases. Keep in mind that in the following argument, we shall use frequently the condition (4.7)–(4.9) and the facts mentioned at the beginning of the proof.

Case 1. $w_{i_1} = uv$

Subcase 1.1. $w_{i_0} = ux$:

By the hypothesis of induction $h_1(w_{i_1}^{-1} \dots) = h_1(v^{-1})$, hence $h_1(w) = h_1(uxv^{-1} \dots) = h_1(u) = h_1(w_{i_0})$.

Subcase 1.2. $w_{i_0} = yv$

Sub-Subcase 1.2.1. $w_{i_2} = yv$:

Since $h_1(w_{i_2} \dots) = h_1(y)$, we have $h_1(w) = h_1(yv(uv)^{-1}y \dots) = h_1(y) = h_1(w_{i_0})$.

Sub-Subcase 1.2.2. $w_{i_2} = ux$:

In this case, either $w_{i_3} = uv$ or $w_{i_3} = yv$, we always have $h_1(w_{i_3}^{-1} \dots) = h_1(v^{-1})$. Thus $h_1(w) = h_1(yv(uv)^{-1}uxv^{-1} \dots) = h_1(yxv^{-1} \dots) = h_1(y) = h_1(w_{i_0})$.

Case 2. $w_{i_1} = yv$

Subcase 2.1. $w_{i_0} = ux$:

$h_1(w_{i_1}^{-1} \dots) = h_1(v^{-1})$ and $h_1(w) = h_1(uxv^{-1} \dots) = h_1(u) = h_1(w_{i_0})$.

Subcase 2.2. $w_{i_0} = uv$:

In this case, either $w_{i_2} = ux$ or $w_{i_2} = uv$, we always have $h_1(w_{i_2} \dots) = h_1(u)$. Thus $h_1(w) = h_1(uv(yv)^{-1}u \dots) = h_1(uy^{-1}u \dots) = h_1(u) = h_1(w_{i_0})$.

Case 3. $w_{i_1} = ux$

Subcase 3.1. $w_{i_0} = uv$:

$h_1(w_{i_1}^{-1} \dots) = h_1(x^{-1})$ and $h_1(w) = h_1(uxv^{-1} \dots) = h_1(u) = h_1(w_{i_0})$.

Subcase 3.2. $w_{i_0} = yv$:

$h_1(w_{i_1}^{-1} \dots) = h_1(x^{-1})$ and $h_1(w) = h_1(yvx^{-1} \dots) = h_1(y) = h_1(w_{i_0})$. $\square$

The above lemma will be used to study substitutions of the form (ux, uv, yv) . The following lemma is the version for (uxv, uv, y) .

Lemma 4.7. *Let u, v, x and y be non-empty words satisfying*

$$h_1(u) \neq h_1(y) \text{ and } h_\infty(v) \neq h_\infty(y);$$

v is not a prefix of xv ;

u is not a suffix of ux .

Let $w_1 = uxv, w_2 = uv, w_3 = y$ and consider the mixed words of the form (4.10). Then

(i) (4.11) holds;

(ii) If $|y| > 1$, (4.12) always holds;

(iii) If $|y| = 1$, (4.12) holds except when $k = 0$, and $w_{i_0} = w_3$.

Proof. (ii) and (iii) are consequences of (i). To show (i), as in the above lemma, we only show $h_1(w) = h_1(w_{i_0}^\epsilon)$ where $\epsilon = +1$.

The cases $k = 0, 1$ are trivial. Let us show the lemma for $k \geq 2$ by induction.

Case 1. $w_{i_0} = uxv$

Subcase 1.1. For all j , $w_{i_j} \neq y$:

Then $w_{i_1} = uv, w_{i_2} = uxv, \dots$

Subsubcase 1.1.1. $k = 2m - 1$:

In this case, $w = ((uxv)(uv)^{-1})^m = ux^m u^{-1}$. Then by Lemma 4.5 we know that $w = +-$ which implies $h_1(w) = h_1(u) = h_1(w_{i_0})$.

Subsubcase 1.1.2. $k = 2m$:

$w = ux^{m+1}v$, the conclusion follows.

Subcase 1.2. Let $w_{i_j} = y$, where $j = \min\{m; w_{i_m} = y\}$

Subsubcase 1.2.1. $j = 2m - 1$:

We have $w = ((uxv)(uv)^{-1})^m y \cdots = ux^m u^{-1} y \cdots$, thus $h_1(w) = h_1(u) = h_1(w_{i_0})$.

Subsubcase 1.2.2. $j = 2m$:

We have $w = ((uxv)(uv)^{-1})^m (uxv)y^{-1} \cdots = ux^{m+1}vy^{-1} \cdots$ then the conclusion follows.

Case 2. $w_{i_0} = uv$. We use a similar proof as the case 1.

Case 3. $w_{i_0} = y$:

For any subcases, it is checked that $h_1(w_{i_1}^{-1} \cdots w_{i_k}^{(-1)^k}) = h_1(v^{-1})$, thus
 $h_1(w) = h_1(yv^{-1} \cdots) = h_1(y) = h_1(w_{i_0})$. $\square$

Lemma 4.8. *Under the notations and conditions of Lemma 4.6 (resp. Lemma 4.7), let $\sigma = (w_1, w_2, w_3)$, then σ is not invertible.*

Proof. Were σ invertible, we would have a Nielsen's cancellation procedure as follows

$$\sigma \rightarrow \cdots \rightarrow (w_1^{(k)}, w_2^{(k)}, w_3^{(k)}) \rightarrow (w_1^{(k+1)}, w_2^{(k+1)}, w_3^{(k+1)}) \cdots \rightarrow I \quad (4.13)$$

where

$$w_j^{(k+1)} \neq \varepsilon \quad (j = 1, 2, 3). \quad (4.14)$$

We will show that:

Claim: $w_i^{(k)}$ ($i = 1, 2, 3$, $k = 1, 2, \dots$) can be written in the form (4.10).

Then the previous lemma implies that $|w_i^{(k)}| \geq 2$ which is in contradiction with (4.13).

Now let us prove the claim by induction.

Suppose that $w_1^{(k)}, w_2^{(k)}, w_3^{(k)}$ are of the form (4.10). Then

$$(w_1^{(k+1)}, w_2^{(k+1)}, w_3^{(k+1)}) = (w_1^{(k)}, w_2^{(k)}, w_3^{(k)}) \circ \tau$$

where $\tau \in \{\pi_1, \pi_2, \phi_l, \phi_r, \iota_1\}$. We shall show that $w_1^{(k+1)}, w_2^{(k+1)}, w_3^{(k+1)}$ are still of the form (4.10).

If $\tau \in \{\pi_1, \pi_2, \iota_1\}$, the conclusion is trivial. So we consider only the case $\tau = \phi_r$. The case for $\tau = \phi_l$ can be treated similarly.

Now $w_1^{(k+1)} = w_1^{(k)} w_2^{(k)}$ (and $w_2^{(k+1)} = w_2^{(k)}$, $w_3^{(k+1)} = w_3^{(k)}$), then by the cancellation condition, we must have $|w_1^{(k)} w_2^{(k)}| \leq |w_1^{(k)}|$ which implies

$$h_\infty(w_1^{(k)}) h_1(w_2^{(k)}) = \varepsilon. \quad (4.15)$$

Without loss of generality we suppose that $h_\infty(w_1^{(k)}) = -$, that is (by the above lemma and the assumption of induction), $w_1^{(k)} = \cdots w_{\lambda_1} w_{\lambda_2}^{-1}$. Then by the cancellation condition (4.15) we must have $w_2^{(k)} = +$, that is, $w_2^{(k)} = w_{\mu_1} w_{\mu_2}^{-1} \cdots$. Consequently $w_1^{(k+1)} = \cdots w_{\lambda_1} w_{\lambda_2}^{-1} w_{\mu_1} w_{\mu_2}^{-1} \cdots$. If $\lambda_2 \neq \mu_1$, the conclusion for $w_1^{(k+1)}$ is proved; if $\lambda_2 = \mu_1$, by cancelling $w_{\lambda_2}^{-1} w_{\mu_1}$ in the above expression and by induction, using the fact that $w_1^{(k+1)} \neq \varepsilon$ (by 4.14), we can show easily that $w_1^{(k+1)}$ is still of the form (4.10). The claim is thus proved. $\square$

The following remarks explain the original idea of the above proof:

Remark 4.1. *There is an essential difference between the proof of the above lemmas and that of Theorem 2.1 in [26] : here w_1, w_2, w_3 are not mixed. The essential point is that, by the assumption of the lemma, we can control the possible cancellations such as $w_1^{-1} w_2 w_3^{-1} \neq - + -$.*

Remark 4.2. *An alternative proof can be done as follows, we omit the details which are not trivial: starting from σ , consider all possible Nielsen's cancellations procedures, we shall get a finite tree of the procedures (which contains several periodic circle), no branch of the tree is reduced to the identity.*

Proposition 4.1. *Suppose that $\sigma = (w_1, w_2, w_3)$ is a non-simple indecomposable substitution. Then we have either $h_1(w_1) = h_1(w_2) = h_1(w_3)$ or $h_\infty(w_1) = h_\infty(w_2) = h_\infty(w_3)$. In other words, w_1, w_2, w_3 must have a common non-empty prefix or suffix.*

Proof. Since σ is indecomposable and invertible, σ satisfies the condition in Corollary 4.1. Hence we can suppose that

Case 1. $\sigma = (w_1, w_2, w_3) = (ux, uv, yv)$

Case 2. $\sigma = (uxv, uv, y)$

Consider the case 1.

If the conclusion of the proposition were not true, we would have that (ux, uv, yv) satisfies the condition of Lemma 4.6 and 4.8, which imply that σ is not invertible, a contradiction.

For the case 2, the proof is the same as the Case 1 except that we apply Lemma 4.7 in stead of Lemma 4.6. $\square$

Lemma 4.9. *Let $z, u, v, x, y \in A^*$. Assume that u, v, x, y satisfy (4.7), (4.8) and (4.9). Denote $w_1 = ux, w_2 = uv, w_3 = yv$,*

$$\begin{aligned}\mathcal{N}_1 &= \{zw_{i_0}w_{i_1}^{-1} \cdots w_{i_{2k-1}}^{-1}z^{-1}; k \geq 1, i_m = 1, 2, 3, \\ &\quad (m = 0, \dots, 2k-1), i_m \neq i_{m+1} (m = 0, \dots, 2k-2)\} \\ \mathcal{N}_2 &= \{zw_{i_0}w_{i_1}^{-1} \cdots w_{i_{2k}}; k \geq 1, i_m = 1, 2, 3, \\ &\quad (m = 0, \dots, 2k), i_m \neq i_{m+1} (m = 0, \dots, 2k-1)\} \\ \mathcal{N}_3 &= \{w_{i_0}^{-1}w_{i_1} \cdots w_{i_{2k-1}}; k \geq 1, i_m = 1, 2, 3, \\ &\quad (m = 0, \dots, 2k-1), i_m \neq i_{m+1} (m = 0, \dots, 2k-2)\} \\ \mathcal{N}_4 &= \{w_{i_0}^{-1}w_{i_1} \cdots w_{i_{2k-1}}w_{i_{2k}}^{-1}z^{-1}; k \geq 1, i_m = 1, 2, 3, \\ &\quad (m = 0, \dots, 2k), i_m \neq i_{m+1} (m = 0, \dots, 2k-1)\} \\ \mathcal{N} &= \mathcal{N}_1 \cup \mathcal{N}_2 \cup \mathcal{N}_3 \cup \mathcal{N}_4\end{aligned}$$

We have

- (1) (4.11) holds;
- (2) If (zw_1, zw_2, zw_3) is cancelled by a Nielsen's procedure to (U_1, U_2, U_3) , then $U_1, U_2, U_3 \in \mathcal{N}$;
- (3) (zux, zuv, zyv) is not invertible.
- (4) (uxz, uvz, yvz) is not invertible.

Proof. (1) is proved in Lemma 4.6. (3) is a consequence of (2). (4) can be proved similarly (by using similar definitions of $\mathcal{N}_k$, or simply by the fact that (4) differs from (3) by an inner automorphism). Let us prove (2) by induction.

Suppose that $\sigma \rightarrow \cdots \rightarrow (V_1, V_2, V_3) \rightarrow (U_1, U_2, U_3)$ where $V_1, V_2, V_3 \in \mathcal{N}$, we show that $U_1, U_2, U_3 \in \mathcal{N}$. With a simple observation of appearances of z and z^{-1} in the expressions above, by the cancellation condition, the conclusion can be proved by using analogous arguments as in the proof of Lemma 4.6. $\square$

Remark 4.3. *When $x, y, u, v, w_1, w_2, w_3$ are given as in Lemma 4.7, similar conclusions of the above lemma hold. The proof is the same.*

Lemma 4.10. *Suppose that $\sigma = (zu_1, zu_2, zu_3)$ (resp. $\sigma = (u_1z, u_2z, u_3z)$) is a non-simple indecomposable substitution where z is a non-empty word. Suppose that u_1, u_2, u_3 have no common prefix and no common suffix. Then there exist a non-trivial invertible substitution σ' , a permutation $\pi \in \mathcal{P}$ and a Fibonacci $f \in \mathcal{F}$, such that*

$$I_{z^{-1}} \circ \sigma = \sigma' \circ f \circ \pi \quad (\text{resp. } I_z \circ \sigma = \sigma' \circ f \circ \pi). \quad (4.16)$$

Proof. Let $w_i = zu_i$ ($i = 1, 2, 3$). First, from the fact that σ is non-simple indecomposable we have

$$w_i \text{ is neither a prefix nor a suffix of } w_j \text{ } (\forall i, j, i \neq j). \quad (4.17)$$

By Lemma 4.2, (w_1, w_2, w_3) is not mixed. It turns out that (u_1, u_2, u_3) is not mixed. By Corollary 4.1, ignoring some permutation, we may write

$$(u_1, u_2, u_3) = (ux, uv, yv) \quad (4.18)$$

$$\text{or } (u_1, u_2, u_3) = (uxv, uv, y). \quad (4.19)$$

We only prove the lemma for the case (4.18). The proof for the case (4.19) is the same, because of Lemma 4.7 and Remark 4.3.

Now we have $\sigma = (zux, zuv, zyv)$. Then $I_{z^{-1}} \circ \sigma = (uxz, uvz, yvz)$. We claim that there exist a non-trivial substitution σ' and a Fibonacci substitution f such that

$$(uxz, uvz, yvz) = \sigma' \circ f \quad (4.20)$$

In fact, if (4.20) fails to hold, denoting $(uxz, uvz, yvz) = (w'_1, w'_2, w'_3)$, we shall have

$$w'_i \text{ is neither a prefix nor a suffix of } w'_j \text{ } (\forall i, j, i \neq j). \quad (4.21)$$

then it follows from (4.17) and (4.21) that u, v, x, y satisfy the condition of Lemma 4.9, which implies that σ is not invertible, a contradiction. Hence $I_{z^{-1}} \circ \sigma$ can be decomposed as in (4.20) and the lemma follows. $\square$

Lemma 4.11. *Let σ be a non-simple indecomposable substitution. Then there exists $z \in \Gamma_A$ such that $I_z \circ \sigma$ is decomposable. That is, there exist non-trivial invertible substitutions σ_1 and σ_2 such that*

$$I_z \circ \sigma = \sigma_1 \circ \sigma_2 \quad (4.22)$$

Furthermore, $|\sigma_i| < |\sigma|$ ($i = 1, 2$).

Proof. Let $\sigma = (w_1, w_2, w_3)$. Since it is indecomposable, for any $i \neq j$, w_i is not a prefix (resp. suffix) of w_j . Hence we may write

$$(w_1, w_2, w_3) = (u_1w, u_2w, u_3w)$$

where w is the maximal common suffix of w_1, w_2, w_3 (if any) and u_1, u_2, u_3 are non-empty words.

Then we have:

$$I_w \circ \sigma = (wu_1, wu_2, wu_3)$$

where u_1, u_2, u_3 have no common suffix.

If (wu_1, wu_2, wu_3) is not indecomposable, the lemma is proved.

Now suppose that (wu_1, wu_2, wu_3) is indecomposable. Since wu_1, wu_2, wu_3 have no common suffix, by Proposition 4.1, we can write

$$I_w \circ \sigma = (wu_1, wu_2, wu_3) = (zu'_1, zu'_2, zu'_3)$$

where $z \neq \varepsilon$ is the maximal common prefix of wu_1, wu_2 , and wu_3 . In this case, u'_1, u'_2, u'_3 have no common prefix and no common suffix. Then by Lemma 4.10, $I_{z^{-1}} \circ I_w \circ \sigma$ is decomposable. That is (by Lemma 4.3) $I_{z^{-1}w} \circ \sigma$ is decomposable.

The last statement of the lemma is an easy consequence of Corollary 4.2. $\square$

Now we are ready to prove our theorems.

Proof of Theorem 3.1:

First we prove Theorem 3.1 (1). Let σ be an invertible substitution. Then there exist $k \geq 1$ and indecomposable substitutions $\sigma_1, \dots, \sigma_k$ such that $\sigma = \sigma_1 \circ \dots \circ \sigma_k$.

If some σ_i is non-simple, then by Lemma 4.11, there exist $z \in \Gamma_A$, invertible substitutions σ'_i and σ''_i such that $\sigma_i = I_{z^{-1}} \circ \sigma'_i \circ \sigma''_i$ and that

$$|\sigma'_i| < |\sigma_i|, \quad |\sigma''_i| < |\sigma_i|. \quad (4.23)$$

Then we repeat such decomposition for σ'_i (resp. σ''_i) and so on. By (4.23), such decomposition will terminate after finite times, that is, finally every factor will be a simple substitution. Hence we can write

$$\sigma = I_{w_1} \circ \tau_1 \circ I_{w_2} \circ \tau_2 \cdots \circ I_{w_n} \circ \tau_n$$

where τ_i ($i = 1, \dots, n$) is a simple substitution and $w_i \in \Gamma_A$ (put $w_i = \varepsilon$ if necessary).

By Lemma 4.3 (and a simple induction), we may write

$$\sigma = I_w \circ \tau_1 \circ \tau_2 \cdots \circ \tau_n$$

where $w \in \Gamma_A$.

Let $\tau = \tau_1 \circ \tau_2 \cdots \circ \tau_n$. It is clear that τ is a simple substitution.

Finally, since $I_{w^{-1}} \circ \sigma = \tau$, Lemma 4.4 implies that $w \in A^*$ or $w^{-1} \in A^*$. Theorem 3.1(1) is thus proved.

For Theorem 3.1(2), denoting

$$|\sigma|_{\min} = \min\{|\sigma(a)|, |\sigma(b)|, |\sigma(c)|\},$$

it is easy to see that it is equivalent to prove the following claim:

Claim: We can choose w in (1) such that $|w| \leq |\sigma|_{\min}$.

Let us prove the claim by induction on $k = |\sigma|$.

If $k \leq 4$ the claim can be verified simply by enumerating all cases.

Suppose that the claim is true for $k \leq n$ (more clearly, for any substitution σ' of length less than n) and that $|\sigma| = n + 1$. By the conclusion (1) of the theorem, there exists $w \in A^*$ or $w^{-1} \in A^*$ such that $I_w \circ \sigma$ is simple. To be specific, we can suppose that $w^{-1} \in A^*$, $\sigma = (w_1, w_2, w_3)$ and that $|w_1| = |\sigma|_{\min}$.

If $|w| \leq |\sigma|_{\min}$, nothing need to prove. Suppose $|w| > |\sigma|_{\min} = |w_1|$. It is easy to see that w_1 is a common prefix of w_1, w_2, w_3 . Let then $w_2 = w_1 w'_2$. We have $(w_1, w_2, w_3) = (w_1, w'_2, w_2) \circ (a, ab, c)$, that is,

$$\sigma = \sigma' \circ f. \quad (4.24)$$

where $\sigma' = (w_1, w_2', w_2)$ is obviously an invertible substitution and $f = (a, ab, c)$ is a Fibonacci substitution.

It is trivial that $|\sigma'| < |\sigma| = n + 1$ and that $|\sigma'|_{\min} \leq |\sigma|_{\min}$, hence by the hypothesis of induction we have the following fact:

There exists $z \in A^*$ (or $z^{-1} \in A^*$) such that $|z| \leq |\sigma'|_{\min}$ and that

$$g := I_z \circ \sigma' \text{ is simple.} \quad (4.25)$$

Then it follows from (4.24) and (4.25) that

$$I_z \circ \sigma = I_z \circ \sigma' \circ f = g \circ f.$$

Since $|z| \leq |\sigma|_{\min}$ and $g \circ f$ is (by definition) simple, the conclusion is proved. $\square$

Proof of Theorem 3.2:

We check easily that if $\sigma = I_w \circ \tau$, then $M_\sigma = M_\tau$. We have also $M_{\sigma \circ \tau} = M_\sigma M_\tau$. The theorem is then a direct consequence of Theorem 3.1. $\square$

Acknowledgement: The authors would like to thank Professors S. Ito, J. Peyrière, and Z.-Y. Wen for helpful discussions.

References

- [1] P. Arnoux et al. *Introduction to finite automata and substitution dynamical systems*, to appear in L. N. in Math., Springer-Verlag.
- [2] P. Arnoux, V. Berthé, H. Ei, S. Ito, *Tilings, quasicrystals, discrete planes, generalized substitutions, and multidimensional continued fractions*. Discrete models: combinatorics, computation, and geometry (Paris, 2001), 059–078 (electronic), Discrete Math. Theor. Comput. Sci. Proc., AA, Maison Inform. Math. Discrèt. (MIMD), Paris, 2001.
- [3] P. Arnoux, S. Ito, *Pisot substitutions and Rauzy fractals*. Journées Montoises d’Informatique Théorique (Marne-la-Vallée, 2000), Bull. Belg. Math. Soc. Simon Stevin 8, no. 2, 181–207, 2001
- [4] F. Axel, D. Gratias, *Beyond Quasicrystals*, Springer-Verlag Berlin Heidelberg, 1995.
- [5] J. Berstel, *Recent results in Sturmian words*, in: *Developments in Language Theory, II (Magdeburg, 1995)*, World Sci. Publishing, River Edge, NJ, 1996, 13–24.
- [6] E. Bombieri and J. Taylor, *Quasicrystals, tilings, and algebraic number theory: some preliminary connections*, Contemporary Mathematics 64 (1987) 241–164.

- [7] A. Cobham, Uniform tag sequences, *Math. System Theory*, 6 (1972) 164-192.
- [8] H. Ei, S. Ito, *Decomposition theorem for invertible substitution*, Osaka J. Math. 34 (1998) 821-834.
- [9] M. Lothaire, *Algebraic combinatorics on words*, To appear in Cambridge University Press, 2002. Available at <http://www-igm.univ-mlv.fr/~berstel/Lothaire>.
- [10] J. M. Luck, C. Godrèche, A. Janner and T. Janssen, *The nature of the atomic surfaces of quasiperiodic self-similar structures*, J. Phys. A: Math. Gen. 26 (1993) 1951–1999.
- [11] R.C. Lyndon, P.E. Schupp, *Combinatorial group theory*, Springer-Verlag, Berlin Heidelberg Newyork, 1977.
- [12] W. Magnus, A. Karrass, D. Solitar, *Combinatorial group theory: Presentations of groups in terms of generators and relations*, Second Revised Edition by Dover Publications, Inc. 1976.
- [13] F. Mignosi et P. Séébold, *Morphisms sturmians et règles de Rauzy*, Journal de Théorie des Nombres de Bordeaux, 5 (1993) 221–233.
- [14] J. Nielsen, *Die Isomorphismen der allgemeinen unendlichen Gruppe mit zwei Erzeugenden*, Math. Ann. 78 (1918) 385–397.
- [15] J. Nielsen, *Die Isomorphismen der freien Gruppen*, Math. Ann. 91 (1924) 169–209.
- [16] J. Peyrière, Z.-X. Wen et Z.-Y. Wen , *Polynomês associées aux endomorphismes de groupes libres*, L’Enseig. Math. t.39 (1993) 153–175.
- [17] Jacques Peyriere, Zhi-Xiong and Zhi-Ying, *Endomorphisms of certain algebras of identities polynomiales*, C. R. Acad. Sci. Paris. t.331, (2000) 111-114.
- [18] M. Queffélec, *Substitution Dynamical System—Spectral Analysis*, Lecture Notes in Mathematics No.1924, New York: Springer-Verlag, 1987.
- [19] Y. Sano, P. Arnoux, S. Ito, *Higher dimensional extensions of substitutions and their dual maps*. J. Anal. Math. 83, 183–206, 2001.
- [20] P. Séébold, *Fibonacci morphisms and Sturmian words*, Theoret.Comput.Sci. 88(1991) 365-384.
- [21] A. Süto, *Singular continues spectrum on a Cantor set of zero Lebesgue measure for the Fibonacci hamiltonian*, J. Stat. Phys. 56 (1989) 527–543.

- [22] Z.-X. Wen and Z.-Y. Wen, *Local isomorphism of the invertible substitutions*, C. R. Acad. Sci. Paris, t. 318, Série I, (1994) 299–304.
- [23] Z.-X. WEN and Z.-Y. WEN, *Some studies of factors of infinite words generated by invertible substitution*, *Formal power series and algebraic combinatorics*, Proc. of 5th Conf. ed. A. Barlotti, M. Delest and R. Pinzani, 1993, p. 455-466.
- [24] Z.-X. Wen and Z.-Y. Wen, *Some properties of the singular words of the Fibonacci word*, European J. Combin. 15(1994) 587-598
- [25] Z.-X. Wen, Z.-Y. Wen, J. Wu, *Invertible substitutions and local isomorphisms*, C. R. Acad. Sci. Paris, Ser. I 334(2002) 629-634.
- [26] Z.X. Wen and Y. Zhang, *Some remarks on invertible substitutions on three letter alphabet*, Chin.Sci.Bulletin, Vol.44 No.19, 1999,1755-1760.